Air Quality Modelling Applications Section Meteorological Service of Canada Environment and Climate Change Canada

REPORT # AQSC-18-031

Documentation for SMOKE-Ready 2015

Air Pollutant Emission Inventory (APEI)

Package version 1

Abstract

This air-quality modelling package includes version 1 of the 2015 Canadian criteria-air-contaminants (CAC) emissions inventory, also referred to as the Air Pollutant Emission Inventory (APEI), in SMOKE-ready format. The APEI is compiled by the Pollutant Inventories and Reporting Division (PIRD) of Environment and Climate Change Canada. The 2015 inventory database compiled by PIRD was then modified by the Air Quality Modelling Applications Section (AQMAS) of Environment and Climate Change Canada for emissions processing with SMOKE for the 2015 Canadian air quality modelling platform. This document describes the delivery contents, including the source-sector-specific SMOKE-ready emission inventory files and all ancillary files needed to generate spatial surrogates and to process the inventory files using the SMOKE emissions processing system.



DOCUMENT HISTORY

| Date | Auteur | Modifications |
|------------|--------------|-----------------------------------|
| 28/08/2018 | Mourad Sassi | First draft |
| 30/10/2018 | Mike Moran | Revisions, added Section 2.2 text |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

Pour des renseignements supplémentaires :

Environnement et Changement climatique Canada Centre de renseignements à la population 7e étage, édifice Fontaine 200, boulevard Sacré-Cœur Gatineau (Québec) K1A 0H3

Téléphone : 1-800-668-6767 (au Canada seulement) ou 819-997-2800

Courriel: ec.enviroinfo.ec@canada.ca



Release: November 2018

Table of Contents

| DOCUMENT HISTORY | 1 |
|---|----|
| List of Acronyms | 3 |
| Delivery Contents | 4 |
| 1 - 2015 Canadian Inventory Description | 5 |
| 1.1 Overview | 5 |
| 1.2 Point source emissions | 5 |
| 1.2.1 NPRI point sources | 5 |
| 1.2.2 UOG point sources | 5 |
| 1.2.3 Wildfire emissions | 6 |
| 1.2.4 Agricultural emissions | 6 |
| 1.3 Area source emissions | 7 |
| 1.3.1 Fugitive particulate matter dust emissions | 7 |
| 1.3.2 Other area sources | 7 |
| 1.4 Transportation sector emissions | 7 |
| 1.4.1 Aircraft emissions | 7 |
| 1.4.2 Marine emissions | 8 |
| 1.4.3 On-road mobile sources | 8 |
| 1.4.4 Off-road mobile sources | 8 |
| 1.4.5 Locomotive emissions | 8 |
| 1.5 Annual total emissions for this 2015 inventory | 9 |
| 2 - Supplementary information | 10 |
| 2.1 Ancillary files | 10 |
| 2.2 Spatial surrogates for processing 2015 Canadian inventory | 10 |
| 3 - References | 13 |
| AOMAS Internal information | 15 |

List of Acronyms

| Acronyms | Description |
|----------|--|
| APEI | Air Pollutant Emission Inventory |
| AQMAS | Air Quality Modelling Applications Section |
| BAU | business as usual |
| CAC | criteria air contaminants |
| СЕРА | Canadian Environmental Protection Act |
| CWFIS | Canadian Wildland Fire Information System |
| ECA | Emission Control Area |
| ECCC | Environment and Climate Change Canada |
| EEA | Economic Exclusion Area |
| E3MC | Energy, Emissions and Economy Model for Canada |
| EPG | Electric Power Generator |
| FEPS | Fire Emission Production Simulator |
| GEM | Global Environmental Multiscale model |
| GHG | Greenhouse Gas |
| I/M | inspection and maintenance |
| LTO | landing and takeoff |
| NAESI | National Agri-Environmental Standards Initiative |
| NAICS | North American Industry Classification System |
| NPRI | National Pollutant Release Inventory |
| ORL | One-record-per-line (SMOKE input format) |
| PEMA | Pollutant Emission Management Area |
| PIRD | Pollutant Inventories and Reporting Division |
| REQA | Réponse aux enjeux en qualité de l'air |
| SCC | Source Classification Code |
| SMOKE | Sparse Matrix Operator Kernel Emissions |
| TF | Transportable Fraction |
| UNECE | United Nations Economic Commission for Europe |
| UOG | upstream oil & gas |

Delivery Contents

| | Archive files included in this delivery: |
|------------------|---|
| # 1 | This .tar file includes the GIS shapefiles used to generate the set of gridded spatial surrogates needed to allocate Canadian 2015 emissions (see details in section2.2). |
| # 2 | Canada_2015_inventory_v1_svn70.tar.gz <ancillary></ancillary> |
| D O C | <documentation></documentation> |
| D O C 2 | <documentation></documentation> |

1 - 2015 Canadian Inventory Description

1.1 Overview

The 2015 Air Pollutant Emissions Inventory (APEI), previously known as the Canadian CAC emissions inventory, was compiled by the Pollutant Inventories and Reporting Division (PIRD) of Environment and Climate Change Canada (ECCC) using both top-down and bottom up approaches (ECCC, 2016). For example, point source emissions were compiled using a bottom-up approach based on the facility level emissions reported to the National Pollutant Release Inventory (NPRI) (EC, 2015), whereas area source emissions were mainly compiled using a top-down approach based on source-specific activity data and emission factors. The PIRD inventory database was further processed by the Air Quality Modelling Applications Section (AQMAS/ECCC) to generate a SMOKE-ready version for the 2015 Canadian air quality modelling platform (Sassi *et al.*, 2016). The source-sector-specific updates included in this delivery are described below in three main sections: point sources (section 1.2); area sources (section 1-3); and all transportation sources (section 1.4).

1.2 Point source emissions

1.2.1 NPRI point sources

The NPRI point source inventory consists of emissions reported by industrial facilities to ECCC and published under the authority of Sections 46–50 of the *Canadian Environmental Protection Act*, (ECCC, 1999). Facilities are required to report emissions of any species for which the annual emissions exceeds a specified threshold. For example, the reporting threshold for SO₂, NO_x, and CO is 20 metric tons/year while it is 10 metric tons/year for VOC species (ECCC, 1999). NPRI facilities are only required to report emissions at their facility level and not by process. Therefore NPRI emissions are only provided with generic SCC codes and more specific SCC values have been assigned by AQMAS to each facility based on its North American Industry Classification System (NAICS) source sub-category. Emissions from some tall smokestacks are separated out with additional VOC species reported.

The air-quality-modelling version of annual NPRI inventory consists of five files (see files #1-5 in

Table **1-1**). The first file contains the emissions for all the species except VOCs. For reference, total VOC emissions are contained in the second file. In order to give the user a choice, two other files containing VOC's emissions for different chemical speciation mechanism, namely CB06 and ADOM-II are included. Finally, one raw VOC inventory consists of detailed individual VOC species and is provided with CAS numbers.

1.2.2 UOG point sources

There are numerous, but relatively small facilities under the Upstream Oil and Gas (UOG) subinventory, including oil and natural gas emissions through exploration, production, and processing to transport to market. Emissions from many UOG facilities are below the NPRI reporting limits (see

Section 1.2.1). Therefore, they are not normally included in the NPRI point source inventory file and their inventory data is compiled separately (see file #6 in

Table 1-1). Specifically excluded are oil sands mining, extraction and upgrading, petroleum refining, and distribution of refined petroleum products, which are included in the NPRI point sources inventory described in section 1.2.1. The UOG emissions has been developed using a bottom-up approach from individual facilities and their associated equipment to the primary emissions sources for fuel combustion, flaring, CO₂ formation and releases, venting (reported and unreported), fugitive and other unintentional releases such as equipment leaks, storage and handling losses, accidental releases and finally, indirect emissions due to fossil-fuel generated electric power consumption. The UOG inventory for year 2015 was projected by PIRD from a 2011 base-year inventory. UOG emissions are provided with SCC codes and facility-level emissions similar to the NPRI, with information for stack parameters for some of the tall stacks.

1.2.3 Wildfire emissions

This is a day-specific wildfire emissions inventory for Canada for the Canadian "fire season" from April 2015 to October 2015. The wildfire emissions estimation is based on the information of active wildfires or "hotspots" provided by the Canadian Wildland Fire Information System (CWFIS, http://cwfis.cfs.nrcan.gc.ca/home), which is operated by Natural Resources Canada (NRCan). The CWFIS detects active wildland fires using the observations from the NASA MODIS and the NOAA/AVHRR satellite-based detection systems. This data file (file #7 in

Table 1-1) contains daily fire locations by latitude and longitude. Environment and Climate Change Canada's GEM model forecasts are then used to determine weather conditions at fire locations and estimate the fuel consumption. The daily total emissions for each fire "hotspot" are calculated based on the BlueSky Fire Emission Production Simulator (FEPS) module for Environment and Climate Canada's FireWork system (Chen *et al.*, 2012). See Pavlovic *et al.* (2016) for more details on the FireWork system. Note also that there is no plume-rise pre-calculations performed for these wildfire emissions. Fire emissions are provided in IDA and EMS-95 format.

Table 1-1. Emission inventory filenames for all point sources.

| Point sources archives files content | Description | |
|---|--------------------------------------|--|
| (1) PS_2015_FF10_noVOC_20181102.csv | NPRI point sources without VOC | |
| (2) <u>PSVOC_2015_FF10_VOC_20181102.csv</u> | NPRI point sources VOC only | |
| (3) <u>PSVOC_2015_FF10_CB6_20181102.csv</u> | NPRI point sources CB06 mechanism | |
| (4) <u>PSVOC_2015_FF10_ADOM_20181102.csv</u> | NPRI point sources ADOM II mechanism | |
| (5) <u>PSVOC_2015_FF10_Species_20181102.csv</u> | NPRI point sources raw VOC species | |
| (6) <u>PS_2015_FF10_UOG.csv</u> | UOG point sources | |
| (7) ptfire_2015_raqdps020fw.tar.gz | Wildfire daily point sources | |
| (8) <u>Airport_LTO_Point_2015_FF10_20180824.csv</u> | Aircraft LTO point sources | |

1.2.4 Agricultural emissions

The 2015 agricultural monthly emissions inventory includes emissions from 56 types of agriculture sources over 274 Canadian census divisions. This inventory data for this sector is compiled as 6 files containing monthly emission values (files #9-14, Table 1-2).

Emissions were provided in point source format as gridded monthly totals for a national grid at 10 km x 10 km resolution. So there is no need to use surrogates to spatially allocate the emissions, but weekly and diurnal temporal profiles are still needed for temporal allocation of monthly emissions to each hour.

Table 1-2. Emission inventory filenames for all agricultural sources.

| Agriculture sources inventory filelist | Description |
|---|-----------------|
| (9) ag animal NH3 2015 FF10 20171214.csv | Animal NH3 |
| (10) ag_animal_VOC_2015_FF10_20171214.csv | Animal VOC |
| (11) ag_havest_2015_FF10_20171214.csv | Harvest PM |
| (12) ag_synth_fertilizer_NH3_2015_FF10_20171214.csv | Fertilizer NH3 |
| (13) ag_tillage_2015_FF10_20171214.csv | Tillage PM |
| (14) ag_winderros_2015_FF10_20171214.csv | Wind erosion PM |

1.3 Area source emissions

1.3.1 Fugitive particulate matter dust emissions

The fugitive dust inventory includes dust emitted from building and road construction, paved and unpaved roads, agriculture, and mine tailings. These area emissions are treated separately, in order to apply Transportable factors (TF) as described by (Pouliot *et al.*, 2010). The data are reported in one file with annual emission value (file #15, Table 1-3). Note that Aeolian dust emissions are not considered.

1.3.2 Other area sources

All other remaining area sources are contained in an emission file, for sectors such as residential, commercial and industrial. All types of fuel combustion, solvents, etc. The data are reported in one file with annual emission value (file #16, Table 1-3).

Table 1-3. Emission inventory filenames for area sources.

| Area sources inventory filelist | Description |
|---|------------------------|
| (15) AS 2015 FF10 20180219 fdust PM.csv | Dust inventory (PM) |
| (16) AS 2015 FF10 20180219 other.csv | All other Area Sources |

1.4 Transportation sector emissions

All transportation emission inventory files are provided separately from point sources or area sources inventory since some are points and others are area sources. Details are given in the subsection 1.4.1 to 1.4.5.

1.4.1 Aircraft emissions

The aircraft sector includes emissions only for Landing/Take-Off (LTO) cycles. Aircraft in-flight emissions and other airport activities are not included. Emissions estimates are based on emission factors for all aircraft and fuel consumption per airport reported annually by Statistics Canada (Report on Energy Supply-Demand). This inventory is reported as monthly point sources (file #8, Table 1-4, also in Table1-1).

1.4.2 Marine emissions

Marine emissions data are provided by the Transportation Division of ECCC. This sector is an aggregation of a number of classes of vessels (1, 2 and 3) that encompass freighters, tankers, tugs, ferries, passenger boats (inboard), fishing boats, and container ships. The marine emissions data are used for all of the Canadian provinces and territories within 200 nautical-miles from the coast referred as the Economic Exclusion Area (EEA). This inventory is reported as annual mobile sources (file #17, Table 1-4).

1.4.3 On-road mobile sources

Emission estimates from on-road vehicles were generated using a bottom-up approach: emission factors were developed for individual vehicle classes and activity data were applied to them. Generally, for non-GHGs, emission factors were developed using the MOVES2014 emissions simulator, originally developed by the EPA, with key datatables modified to reflect Canadian (provincial/territorial/regional) conditions.

This inventory is compiled as monthly emissions values for different vehicle types. It is aggregated at the provincial level except for two provinces where inspection/maintenance (I/M) programs or other sub-provincial information are available. The inventory for these two provinces was split into sub-regions: British Columbia into two sub-regions and Ontario into four sub-regions. Five road types were also used to improve the spatial allocation (files #18-19, Table 1-4).

1.4.4 Off-road mobile sources

Emissions estimates for all off-road vehicles and engines are based on the EPA's NONROAD v6.7 tool. This inventory is compiled as monthly emissions values from different types of engines and fuels. This inventory is at the provincial level except for four provinces where sub-provincial information is available: British Columbia has two sub-regions, Alberta has two sub-regions, Ontario has four sub-regions, and Quebec has two sub-regions (file #20, Table 1-4).

1.4.5 Locomotive emissions

Rail emissions were obtained from the Transportation Division of ECCC. This inventory is reported as annual mobile sources (file #21, Table 1-4).

REQA/AQMAS 8 Release: November 2018

Table 1-4. Emission inventory filenames for the transportation sector.

| Transportation sources inventory filelist | Description | |
|--|-----------------------------|--|
| (8) <u>Airport_LTO_Point_2015_FF10_20180824.csv</u> | Aircraft LTO inventory | |
| (17) Marine_2015_FF10_20180824.csv | Marine inventory | |
| (18) OnRoad 2015 FF10 20180911.csv (19) OnRoad Refueling 2015 FF10 20180914.csv | On-road mobile & refueling | |
| (20) NonRoad 2015_FF10_20180914.csv | Off-road engine & equipment | |
| (21) Rail_2015_FF10_20180824.csv | Locomotive | |

1.5 Annual total emissions for this 2015 inventory

Emissions from all of the emission inventory files described in the previous sections (1.2 to 1.4) are included in the summary table (Table 1-5) below with their annual total emission for reference.

Table 1-5. Canadian air pollutant annual emissions for 2015 (From APEI14_2015).

| 2015 emissions in tonnes | СО | NH3 | NOX | PM10 | PM2_5 | SO2 | VOC |
|--------------------------|-----------|---------|-----------|-----------|-----------|-----------|-----------|
| Industrial Sources | 1,190,152 | 15,184 | 622,752 | 151,725 | 61,838 | 765,121 | 816,730 |
| Non Industrial Sources | 1,265,302 | 2,881 | 233,428 | 175,058 | 171,243 | 259,996 | 235,389 |
| Miscellaneous Sources | 9,737 | 712 | 76 | 18,182 | 16,745 | 88 | 378,510 |
| Off-road mobile Sources | 1,397,071 | 651 | 570,038 | 27,409 | 26,269 | 11,443 | 182,187 |
| On-road mobile sources | 1,573,028 | 6,486 | 389,053 | 23,988 | 13,484 | 1,461 | 125,879 |
| Open Sources | 147,490 | 459,280 | 11,320 | 6,851,070 | 1,330,700 | 10,709 | 116,434 |
| Total w/o wildfires | 5,582,779 | 485,195 | 1,826,666 | 7,247,433 | 1,620,280 | 1,048,819 | 1,855,129 |

2 - Supplementary information

This section describes other important information also included with the package.

2.1 Ancillary files

SMOKE ancillary files, such as the temporal cross-reference file and temporal profile libraries, spatial cross-reference file, costcy and pstk are included in the ancillary directory of the archive. The spatial cross-reference file "amgref.CAN_T3_T4_PRG2013.txt" should be used for processing on-road and off-road mobile source emissions, while the other spatial cross-reference file, "amgref.CAN2006_noNAESI_12Feb2018.txt", is for processing emissions from all other non-point source emissions.

2.2 Spatial surrogates for processing 2015 Canadian inventory

A set of GIS shapefiles that are used to build the set of Canadian 2015 spatial surrogate fields for Canadian emissions processing is included in the package along with control files needed by the CMAS Surrogate Allocator package (www.cmascenter.org/sa-tools/). All of these files are contained in the compressed tar file named "2015CA_Inv_Package_surrogates.tar.gz".

2.2.1 Updates to the Canadian spatial surrogate database

Since the release of the SMOKE-ready 2013 Canadian APEI package in 2017, some new GIS shapefiles and companion spatial surrogate fields have been developed and added to this package:

- 2016 Census population and dwelling data is used to generate surrogates 100–104, to replace 2006 Census data (more information about the capped population surrogates may be found in Zheng et al., 2015);
- New airport shapefile with detailed runway geometry is used to generate surrogates 901–905;
- Updated Athabasca Oil Sand shapefile is used to generate surrogate 1255;
- Updated residential wood combustion shapefile for surrogate 951, using 2011 household wood consumption data for 13 provinces and 28 Census Metropolitan Areas. The wood consumption was spatially distributed into Dissemination Areas by dwelling counts, urban/rural, the latitude, and the distances to forest (Zheng et al., 2015);
- NAESI surrogates have been removed because agriculture emissions are now reported in point source format at 10 km resolution
- More information is available in the package in files
 "surrogate_generation_canada_Aug2018.csv", "shapefile_catalog_CA_Aug2018.csv",
 "surrogate_ID_CA_Aug2018.csv", and "surrogate_specification_CA_Aug2018.csv".

2.2.2 Summary of Canadian shapefiles used for spatial surrogate generation

The following table provides a summary of the GIS shapefiles and related spatial surrogates grouped by type of surrogate.

| Catalogue | Shapefiles | Surrogates |
|----------------|---|----------------|
| Population | gpr_gda_000a16a_e_sphere_3Nov2017.shp | 100, 101, 102, |
| and Dwelling | gpr_gda_000b11a_SimpP_100m_water_Dissolve.shp | 103, 104, 105 |
| | gda_000b11a_e_SimpPoly_MaxOffset_100m_pop_dwell_Jul2014.shp | |
| | gda_000b16a_e_PopDwell_SimpPoly_100m_noLakes_sphere_1Nov201 | |
| | 7.shp | |
| On-road | gcd_000b11a_e_CDID_BC2_ON4_Simpl10m_water.shp | 200, 210, 220, |
| | NRN_CA_Simp2_16Apr2016_sphere.shp | 230, 240 |
| | gda_000b16a_e_PopDwell_SimpPoly_100m_noLakes_sphere_1Nov201 | |
| | 7.shp | |
| Off-road T4 | gcd_000b11a_e_CDID_BC2_Que2_ON4_Simpl10m_water.shp | 1251, 1252, |
| | da2006_pop_labour_SimplifyP_MaxOffset-100m_sphere_noLakes.shp | 1253, 1254, |
| | allroads.shp | 1255, 1256, |
| | construction_other.shp | 1257, 1258, |
| | mine.shp | 1259, 1260, |
| | naesi_fert.shp | 1261, 1262, |
| | unpaved_ur.shp | 1263, 1264, |
| | wood_industries.shp | 1265 |
| | OS_MinePit_D_v2.shp | |
| | lu100.shp | |
| | othjet_II.shp | |
| | CANRAIL.shp | |
| Airport, rail, | Airport_movements_2006_MultiRingBuffer.shp | 901, 902, 903, |
| and marine | Aviation_Extended-Runways-including-runways-with-buffer- | 904, 905, 932, |
| | Spring2017.shp | 945 |
| | CANRAIL.shp | |
| | Off-Road-using-OSM-Apron-of-155-Aiport-Locations-with-Traffic-data- | |
| | by-Province-STATCAN-2015-Spring2017.shp | |
| | Surrogate-2017-matches-Inventaire-2013-by-Keys-with-FAKEFIPS-6- | |
| | polygon.shp | |
| | marine.shp | |
| Land cover | CONT42_pop_water_Clip_b.shp | 350, 948 |
| | coast_inland_water_Union_BND_m.shp | |
| 0.1 | treesa.shp | 100 111 110 |
| Others: | da2006_pop_labour_SimplifyP_MaxOffset-100m_sphere_noLakes.shp | 106, 111, 113, |
| industrial, | gda_000b11a_e_SimpP_100m_WoodConsump_1Aug2014.shp | 116, 211, 212, |
| commercial, | gpr_gda_000b11a_SimpP_100m_water_Dissolve.shp | 221, 222, 233, |
| agricultural, | lowmedjet_ll.shp | 308, 321, 323, |
| and fugitive | naesi_fert.shp | 324, 326, 327, |
| open | naesi_livestk.shp | 331, 412, 448, |
| sources | othjet_ll.shp | 562, 921, 923, |
| | prov2006_Jan2012_SimplifyPo_MaxOffset-100m_water_sphere.shp | 924, 925, 926, |
| | treesa.shp | 927, 940, 942, |

REQA/AQMAS 11 Release: November 2018

2015 APEI Package-version 1

AQSC-18-031 (rev 70)

| ua2001.shp | 946, 949, 951, |
|----------------|----------------|
| unpaved4.shp | 955, 960, 970, |
| unpaved5.shp | 980, 990, 996 |
| unpaved_ur.shp | |

REQA/AQMAS 12 Release: November 2018

3 - References

Chen, J., R. Pavlovic, A. Pankratz, and K. Anderson, 2012. Evaluation of wildfire emissions in the Canadian GEM-MACH air quality forecast system. *20th Emissions Inventory Conference*, 13-16 Aug., Tampa, Florida. https://www3.epa.gov/ttn/chief/conference/ei20/session2/jchen_pres.pdf

Environment Canada, 2015. Overview of Reviewed Facility-Reported Data National Pollutant Release Inventory (NPRI) 2015. (Cat. No.: En81-24-E-PDF ISSN: 2369-7733) https://www.ec.gc.ca/inrp-npri/default.asp?lang=En&n=386BAB5A-1&printfullpage=true&wbdisable=false

Environment and Climate Change Canada (1999). The Canadian Environmental Protection Act. http://www.ec.gc.ca/lcpe-cepa/default.asp?lang=En&n=26A03BFA-1.

Environment and Climate Change Canada, 2016. Canadian Environmental Sustainability Indicators: Air Pollutant Emissions. (Cat. No.: En4-144/22-2016E-PDF ISBN: 978-0-660-04749-2). http://www.ec.gc.ca/indicateurs-indicators/default.asp?lang=en&n=E79F4C12-1

Pavlovic, R., J. Chen, K. Anderson, M.D. Moran, P.-A. Beaulieu, D. Davignon and S. Cousineau, 2016. The FireWork air quality forecast system with near-real-time biomass burning emissions: Recent developments and evaluation of performance for the 2015 North American wildfire season. *J. Air & Waste Manage. Assoc.*, http://doi.org/10.1080/10962247.2016.1158214.

Pouliot et al., 2010. Assessing the Anthropogenic Fugitive Dust Emission Inventory and Temporal Allocation Using an Updated Speciation of Particulate Matter. From book Air Pollution Modeling and its Application XXI (pp.585-589).

https://www.researchgate.net/publication/226521498 Assessing the Anthropogenic Fugitive D ust Emission Inventory and Temporal Allocation Using an Updated Speciation of Particulat e Matter

Sassi, M., S. Cousineau, A. Duhamel, S. Ménard, J. Racine, C. Zaganescu and R. Mashayekhi. The 2013 Canadian air quality modelling platform and the base future cases used for policy regulations. 15th Annual CMAS Conference, Chapel Hill, NC, October 2016 https://www.cmascenter.org/conference/2016/abstracts/sassi 2013 canadian 2016.pdf

Zheng, Q., M.D. Moran, J. Zhang, E. Tanguay, D. Niemi, and M. Sassi, 2015. Development of some improved Canadian spatial surrogates. Poster, 21st International Emissions Inventory Conference, 13-17 April, San Diego. [see http://www.epa.gov/ttn/chief/conference/ei21/poster/qzheng.pdf]

Contact

For any request regarding this delivery, please send an email to the distribution list EC.REQA.EC@canada.ca. A member of the team will get back to you as soon as possible.

AQMAS Internal information

Historical, revision information and local path locations

Canada 2015 inventory v1 svn70.tar.gz

Final package prepared by Mourad Sassi and Rabab Mashayekhi (November 2018) based on: https://ulysse.cmc.ec.gc.ca/svn/emission_inventory/2015 (revision #70, November 2nd 2018)

Local extraction path:

/fs/cetus/fs4/aq/afsusno/Emissions/EmissionsInventoryPackage

Local delivery package path:

/fs/cetus/fs4/aq/afsusno/Emissions/EmissionsInventoryPackage/Canada_2015_inventory_v1_svn70

All related information regarding this request will be recorded under Mantis issue #2805